

3-Branes on Eguchi-Hanson $6D$ Instantons

Harvendra Singh

Saha Institute of Nuclear Physics
1/AF Bidhannagar, Kolkata 700064, India

E-mail: h.singh[AT]saha.ac.in

Abstract

We use the approach used by Eguchi-Hanson in constructing four-dimensional instanton metrics and construct a class of regular six-dimensional instantons which are nothing but $S^2 \times S^2$ resolved conifolds. We then also obtain D3-brane solutions on these EH-resolved conifolds.

1 Introduction

The AdS/CFT conjecture [1, 2, 3] relates string theory on anti-de Sitter bulk spacetime to a four-dimensional conformal gauge theory on the boundary. It serves as being one of the concrete realisation of the holographic idea [4]. In this picture the $SU(N)$ super-Yang-Mills theory lives on the boundary of the bulk $AdS_5 \times S^5$ space. Since then this gauge-gravity duality has been extensively tested and has provided a fruitful alternative in understanding of various aspects of gauge theories, both qualitatively and quantitatively. The large body of work is cited in some reviews, for example [5].

It has been rather interesting to study AdS/CFT for the class of manifolds $AdS_5 \times T^{1,1}$ which have less supersymmetry, where $T^{1,1}$ is an Einstein space. This requires constructing D3-brane solutions over $M_4 \times Y^{p,q}$ spaces where $Y^{p,q}$ is the six-dimensional Calabi-Yau cone. These cone-like geometries are singular at the tip of the cone, but deformations on these conifold geometries can be performed [6] so that they become regular Calabi-Yau geometries. Several new resolved solutions have recently appeared in literature [7, 8].

In this short note we first construct a regular Calabi-Yau cone solution which we obtain by adopting standard method of constructing Eguchi-Hanson 4D instantons. We call them EH-resolved cone as they share some unique properties with Eguchi-Hanson instantons. But there are standard techniques to resolve these cones, see [6, 9]. We then look for D3-brane solutions on these EH-resolved cones. The paper is organised as follows. In the section-2, we review Eguchi-Hanson 4D instanton geometry. In section-3 we construct six-dimensional regular cone-like geometry using Eguchi-Hanson ansatz. We then obtain D3-brane configurations over these spaces in the section-4. The results are summarised in the last section.

2 The 4D EH instantons

The Euclidean gravity solutions with finite action and a self-dual curvature are manifolds that are generally classified as gravitational instantons. The gravitational fields are localised in space and the metric becomes asymptotically locally flat at infinity.¹ In this section we review the main aspects of the Eguchi-Hanson 4D solutions.

In the case of the Eguchi-Hanson (EH) manifold the metric ansatz is [12]

$$ds_{EH}^2 = \frac{dr^2}{g(r)} + \frac{r^2}{4}(d\theta^2 + \sin^2 \theta d\phi^2) + \frac{r^2}{4}g(r)(d\psi + \cos \theta d\phi)^2 \quad (1)$$

where $r^2 = t^2 + x^2 + y^2 + z^2$. The t, x, y, z are Cartesian coordinates which define the base space which is R^4 . There is a complete radial symmetry and one is free to pick

¹It has been recently found that the information-geometry on the solution space of these instantons has a constant negative curvature [13]. The results are found to be quite similar to the information-metric case of Yang-Mills instantons [14].

the center in the base space. The above metric solves Euclidean gravity equations in flat space provided $g(r)$ satisfies a monopole like first order equation

$$g'(r) + \frac{4}{r}(g - 1) = 0 \quad (2)$$

which has an immediate solution [12]

$$g(r) = 1 - \frac{a^4}{r^4}, \quad (r \geq a) \quad (3)$$

where a is an integration constant. By redefining the coordinate $\rho^4 = r^4 - a^4$ one can also express metric as [12]

$$ds_{EH}^2 = \left(1 + \frac{a^4}{\rho^4}\right)^{-\frac{1}{2}} (d\rho^2 + (\rho^2/4)(d\psi + \cos\theta d\phi)^2) + \left(1 + \frac{a^4}{\rho^4}\right)^{\frac{1}{2}} \frac{\rho^2}{4} (d\theta^2 + \sin^2\theta d\phi^2) \quad (4)$$

with the coordinate ranges

$$0 \leq \rho \leq \infty, \quad 0 \leq \theta \leq \pi, \quad 0 \leq \phi \leq 2\pi, \quad 0 \leq \psi \leq 2\pi. \quad (5)$$

A non-singular Maxwell field with self-dual field strength can be introduced in the Eguchi-Hanson background

$$\begin{aligned} A_{(1)} &= \frac{a^2}{\sqrt{\rho^4 + a^4}} \sigma_z, \\ F_{(2)} &= \frac{2a^2}{\rho^4 + a^4} (e^3 \wedge e^0 + e^1 \wedge e^2). \end{aligned} \quad (6)$$

Thus for EH instanton,

$$(F_{\mu\nu})^2 = \frac{16a^4}{(\rho^4 + a^4)^2}. \quad (7)$$

Though a is an integration constant, but it is related to the presence of Maxwell field here. For the instantons located away from the origin in R^4 , one can introduce 4 new parameters, x_0^i , as the position variables. This is possible because the EH metric and the gauge background have complete radial symmetry, so $\rho^2 = |x - x_0|^2$. However, in any case the center coordinates, x_0^i , of the EH metric (4) and the gauge field (7) must coincide in order that the solution exists. It is interesting to note that the Eq.(7) is strikingly similar to the Yang-Mills instanton field strength in [10, 11, 14], the only difference being the distribution of powers in the denominator. This fact did lead us to conclude that the information metric approach [14] to holography can also be studied for this class of curved EH instantons also; see for more details [13].

3 A EH-resolved 6D conifold

We construct a class of regular six-manifolds which are the Ricci-flat Kähler solutions of the Einstein equations. For this purpose, we know the existence of the conifold geometries

[6, 15] and corresponding *AdS/CFT* analysis has been studied in detail by [7, 8]. The many deformed and resolved conifolds have also been worked out in [8, 9] previously. As we are interested in Eguchi-Hanson class of instantons on the conifold, so we look for a metric ansatz resembling the Eguchi-Hanson $4D$ metric

$$ds_6^2 = \frac{dr^2}{f(r)} + \frac{r^2}{6} \left((d\theta_1^2 + \sin^2 \theta_1 d\phi_1^2) + (d\theta_2^2 + \sin^2 \theta_2 d\phi_2^2) \right) + \frac{r^2}{9} f(r) (d\psi + \sum_{i=1}^2 \cos \theta_i d\phi_i)^2. \quad (8)$$

The angular coordinate ranges are taken as

$$0 \leq \theta_i \leq \pi, \quad 0 \leq \phi_i \leq 2\pi, \quad 0 \leq \psi \leq 2\pi. \quad (9)$$

One will easily note that the $r = \text{constant}$ sections of the above metric are indeed $T^{1,1}$ geometry, which is an Einstein space with positive curvature. Now all the $R_{\mu\nu} = 0$ equations are solved by the metric (8) provided $f(r)$ satisfies the following monopole type equation,

$$f' + \frac{6}{r}(f - 1) = 0. \quad (10)$$

Note that this first order equation appears in various instanton solutions including the Yang-Mills and the $4D$ Eguchi-Hanson one. These are consequences of BPS conditions. The above differential equation has one trivial solution $f = 1$, in which case the metric becomes that of the singular Calabi-Yau cone over the base $T^{1,1}$ [6]

$$ds_6^2 = dr^2 + r^2 ds_{T^{1,1}}^2 \quad (11)$$

which is Ricci-flat and the Kähler 2-form is given by

$$J = \frac{r^2}{6} (\sin \theta_1 d\theta_1 \wedge d\phi_1 + \sin \theta_2 d\theta_2 \wedge d\phi_2) + \frac{r}{3} (d\psi + \cos \theta_i d\phi_i) \wedge dr. \quad (12)$$

The nontrivial solution of (10) is

$$f(r) = 1 - \frac{a^6}{r^6}, \quad a \leq r < \infty \quad (13)$$

where a is an integration constant. We can take it to be the size of the instantons, or the measure of deformation.² For the solution (13) the metric becomes

$$ds_6^2 = \frac{dr^2}{1 - \frac{a^6}{r^6}} + \frac{r^2}{6} \left((d\theta_1^2 + \sin^2 \theta_1 d\phi_1^2) + (d\theta_2^2 + \sin^2 \theta_2 d\phi_2^2) \right) + \frac{r^2}{9} \left(1 - \frac{a^6}{r^6} \right) (d\psi + \cos \theta_i d\phi_i)^2. \quad (14)$$

which appears quite the same as Eguchi-Hanson metric (1). This EH-resolved six-manifold is topologically $R^+ \times S^2 \times S^3$. An interesting *unique* property of the EH-resolved metric, that is why we call it so, is that the determinant of the resolved metric remains the same

² The third solution with plus sign $f(r) = 1 + \frac{a^6}{r^6}$ will have a naked singularity at $r = 0$.

as the unresolved metric. It was also the case with the 4D instantons. The apparent coordinate singularity at $r = a$ is removable, it can be seen as follows. Near $r = a$ we define new radial coordinate $u^2 = r^2(1 - \frac{a^6}{r^6})$, so the metric (14) can be written as

$$ds_6^2 = \frac{du^2}{(1 + \frac{2a^6}{r^6})^2} + \frac{u^2}{9}(d\psi + \cos\theta_i d\phi_i)^2 + \frac{r^2}{6} \left((d\theta_1^2 + \sin^2\theta_1 d\phi_1^2) + (d\theta_2^2 + \sin^2\theta_2 d\phi_2^2) \right). \quad (15)$$

So near $r = a$ it becomes

$$ds_6^2 \approx \frac{1}{9}(du^2 + u^2(d\psi + \cos\theta_i d\phi_i)^2) + \frac{a^2}{6} \left((d\theta_1^2 + \sin^2\theta_1 d\phi_1^2) + (d\theta_2^2 + \sin^2\theta_2 d\phi_2^2) \right), \quad (16)$$

and thus the metric can be made regular with the topology being $R^2 \times S^2 \times S^2$ near $r = a$, if the range of ψ is restricted to $0 \leq \psi \leq 2\pi$. While for $r \gg a$ the metric (14) becomes the metric over the Kähler cone (11).

In any case the Kähler 2-form for the metric (14) is ³

$$J = e^5 \wedge e^0 + e^1 \wedge e^2 + e^3 \wedge e^4 \quad (17)$$

The vielbeins are

$$\begin{aligned} e^0 &= (1 - \frac{a^6}{r^6})^{-1/2} dr, \quad e^1 = \frac{r}{\sqrt{6}} d\theta_1, \quad e^2 = \frac{r}{\sqrt{6}} \sin\theta_1 d\phi_1, \quad e^3 = \frac{r}{\sqrt{6}} d\theta_2, \\ e^4 &= \frac{r}{\sqrt{6}} \sin\theta_2 d\phi_2, \quad e^5 = \frac{r}{3} (1 - \frac{a^6}{r^6})^{1/2} (d\psi + \cos\theta_1 d\phi_1 + \cos\theta_2 d\phi_2) \end{aligned} \quad (18)$$

3.1 Adding the tensor fields

We now wish to introduce a pair of second rank tensor fields $B_{\mu\nu}$ and $C_{\mu\nu}$ in the 6-dimensional manifold (14). The most plausible combined action is

$$\int \left(R * 1 - \frac{1}{2!} H_{(3)} \wedge * H_{(3)} - \frac{1}{2!} F_{(3)} \wedge * F_{(3)} - \frac{1}{2!} F_{(1)} \wedge * F_{(1)} + \text{boundary} - \text{terms} \right) \quad (19)$$

where field strengths are given by $H_{(3)} = dB_2$ and $F_{(3)} = dC_2 - \chi dB_2$ and $F_{(1)} = d\chi$. In our Hodge-dual convention: $*1 = e^1 \wedge \dots \wedge e^6 \equiv \sqrt{g}[d^6x]$. The field equations which follow from the above action are

$$\begin{aligned} d * F_3 &= 0, \quad d(*H_3 - \chi * F_3) = 0, \\ d * d\chi &= dB_2 \wedge * dC_2 \end{aligned} \quad (20)$$

Thus a vanishing χ solution exists provided we ensure $F_{\mu\nu\lambda} H^{\mu\nu\lambda} = 0$. For $\chi = 0$ the tensor field equations are readily solved provided we take

$$* F_3 = -H_3, \quad * H_3 = F_3. \quad (21)$$

From here, one can construct a complex harmonic $(2, 1)$ -form $G_3 = F_3 + iH_3$ which satisfies the following complex self-duality relation

$$* G_3 = iG_3. \quad (22)$$

and its energy-momentum tensor identically vanishes.

³ Note that it can be written as $J \equiv dA = d(-\frac{r^2}{6}(d\psi + \cos\theta_i d\phi_i))$. Also $J \wedge J \wedge J \equiv \text{Vol}[M_6]$.

3.2 The singular B -field solution

We now construct a solution of the coupled field equations of Einstein and 2-rank tensor fields. It is found that a solution exists where the metric is taken to be the Eq.(14) and the B field as

$$B_2 = \frac{m}{2} \log \frac{r^6 - a^6}{r_0^6} \omega_2, \quad (23)$$

where ω_2 is a closed 2-form over S^2 's

$$\omega_2 = \frac{1}{2}(\sin \theta_1 d\theta_1 \wedge d\phi_1 - \sin \theta_2 d\theta_2 \wedge d\phi_2).$$

Then the corresponding 3-form field strength becomes

$$H_{(3)} = \frac{3mr^5}{(r^6 - a^6)} dr \wedge \omega_2 \quad (24)$$

where m is a constant parameter. The field strength $F_{(3)}$ is determined by the Hodge-dual relations (21) and it is

$$F_{(3)} = m \omega_3 \quad (25)$$

where $\omega_3 = (d\psi + \cos \theta_i d\phi_i) \wedge \omega_2$. There is a constant flux through S^3

$$\int_{S^3} F_3 = m \quad (26)$$

which will be quantized. Due to the self-dual property (22) the field strength G_3 is harmonic and the corresponding Euclidean energy-momentum tensor identically vanishes. This has been explicitly checked by us. This is the reason that the Einstein equations retain their empty-space form $R_{\mu\nu} = 0$. Also we have

$$F_{\mu\nu\lambda} H^{\mu\nu\lambda} = 0$$

for this background which is required for axion to be vanishing.

However, the field expression

$$\frac{1}{2 \cdot 3!} (H_{\mu\nu\lambda})^2 = \frac{(9m)^2}{(r^6 - a^6)} \quad (27)$$

is singular at $r = a$. Since $(F)^2 = (H)^2 \sim \frac{1}{r^6}$ for large r , the asymptotic behavior of these B -instantons is different from that of the regular $4D$ instantons discussed in the last section. Note that, the tensor-fields remain divergent at $r = a$ even though the metric (14) is regular there. The action for the tensor fields diverges when integrated over the range $a \leq r \leq \infty$ because of the singularity of the tensor fields at $r = a$. So this cannot represent a regular instanton background unlike Eguchi-Hanson background. This singularity at $r = a$ is similar to the Klebanov-Strassler conifold singularity where the fields are singular at the tip of the cone $r = 0$. Whole of this structure reduces to the singular conifold case when $a = 0$. One had to deform the conifold in order to avoid the singularity [7].

4 Embedding in ten dimensions

The six-dimensional conifolds can also be embedded in ten-dimensional IIB string theory quite elegantly [7]. It is of main interest to insert D3-branes on the above resolved conifold (14). The full ten-dimensional ansatze are then

$$\begin{aligned}
ds^2 &= h(r)^{-\frac{1}{2}}(-dt^2 + dx \cdot dx) + h(r)^{\frac{1}{2}} \left(\frac{dr^2}{f(r)} + \frac{r^2}{9} f(r) (\tilde{\psi})^2 + \frac{r^2}{6} \sum_{i=1}^2 (d\theta_i^2 + \sin^2 \theta_i d\phi_i^2) \right) \\
F_5 &= dh^{-1} \wedge dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3 + Q(r) \omega_2 \wedge \omega_3 \\
\Phi &= 0, \quad \chi = 0 \\
F_3 &= m\omega_3, \quad B_2 = \frac{m}{2} \log \frac{r^6 - a^6}{r_0^6} \omega_2
\end{aligned} \tag{28}$$

where $f(r) = 1 - \frac{a^6}{r^6}$ and we defined $\tilde{\psi} \equiv d\psi + \sum_i \cos \theta_i d\phi_i$. We have to ensure that field strength $F_5 \equiv dC_4 + B_2 \wedge F_3$ satisfies its equations of motion

$$dF_5 = d *_5 F_5 = H_3 \wedge F_3$$

and is also self-dual. Thus all field equations are solved provided

$$\begin{aligned}
M(r) &= \frac{m^2}{2} \log \frac{r^6 - a^6}{r_0^6} \\
fr^5 \partial_r h &= -Q(r) = -M(r) - c_0
\end{aligned} \tag{29}$$

where c_0, r_0 are constants. An exact solution exists when $a = 0$ (unresolved case)

$$h(r) = b_0 + \frac{1}{r^4} \left(\frac{c_0}{4} + \frac{3m^2}{4} \log\left(\frac{r}{r_0}\right) + \frac{3m^2}{16} \right) \tag{30}$$

which is the solution of [7] provided we set $c_0 = 16\pi g_s N$. In the absence of 3-form flux, $m = 0$, the above solution represents N D3-branes on a singular conifold.

When $a \neq 0$, an exact solution is a bit cumbersome. We can however make following suitable radial coordinate choice,

$$\rho^6 = r^6 - a^6, \quad 1 - \frac{a^6}{r^6} = \left(\frac{\rho}{r}\right)^6, \quad 1 + \frac{a^6}{\rho^6} = \left(\frac{r}{\rho}\right)^6 \tag{31}$$

so that the EH-resolved metric (14) in the new coordinates becomes

$$ds_6^2 = \frac{d\rho^2 + \frac{\rho^2}{9} (\tilde{\psi})^2}{q(\rho)^{2/3}} + \frac{\rho^2}{6} q(\rho)^{1/3} \sum_{i=1}^2 (d\theta_i^2 + \sin^2 \theta_i d\phi_i^2) \tag{32}$$

with the function $q(\rho) = 1 + a^6/\rho^6$ and the new coordinate range $0 \leq \rho \leq \infty$. The singular conifold is obtained whenever we set $a = 0$. In these coordinates 10-dimensional metric (28) will look like

$$ds^2 = h(\rho)^{-\frac{1}{2}} (dx_\mu^2) + h(\rho)^{\frac{1}{2}} \left(\frac{d\rho^2 + \frac{\rho^2}{9} (\tilde{\psi})^2}{q(\rho)^{\frac{2}{3}}} + \frac{\rho^2}{6} q(\rho)^{\frac{1}{3}} \sum_{i=1}^2 (d\theta_i^2 + \sin^2 \theta_i d\phi_i^2) \right) \tag{33}$$

We find that the solution of the equation (30) in the region $\rho \gg a$ can be expressed as

$$h(\rho) = b_0 + \frac{4\pi g_s N}{\rho^4} \left(1 + \frac{4}{15} \frac{a^6}{\rho^6} + \dots \right) + \frac{3m^2}{\rho^4} \left(\frac{1}{16} \log\left(\frac{10\rho^4}{\rho_0^4}\right) + \frac{1}{150} \frac{a^6}{\rho^6} \log\left(\frac{10\rho^{10}}{\rho_0^{10}}\right) + \dots \right) \quad (34)$$

which gives exactly the metric in [7] once $a = 0$. But the conifold is the resolved conifold now. Even in the absence of H-flux ($m = 0$) the brane geometry does not exactly look like $AdS_5 \times T^{1,1}$ if b_0 is dropped, that is when $4\pi g_s N \gg \rho$. The AdS space will be obtained only when we take $4\pi g_s N \gg \rho \gg a$, that is the branes must be located far away from the origin of the conifold. Thus if large number of coincident D3-branes are placed in the region $\rho \gg a$, the near horizon geometry would become more or less $AdS_5 \times T^{1,1}$. That could be seen by dropping the constant b_0 in (34) and setting $m = 0$

$$h(\rho) \sim \frac{4\pi g_s N}{\rho^4} \left(1 + \mathcal{O}\left(\frac{a^6}{\rho^6}\right) + \dots \right) \quad (35)$$

When $m \neq 0$ the same will apply but instead there will be flux dependent deformations, which are logarithmic, of the AdS_5 geometry with interesting consequences in CFT. The D3-Branes on resolved conifolds have several interesting results in the dual CFT [7].

5 Summary

In this short note we constructed regular six-dimensional conifolds simply starting from Eguchi-Hanson like ansatz and solving the resultant first order monopole like equations. We then switched on F_3 flux, the field strengths are found to be singular even for the resolved case. We then construct solutions in ten-dimensions and obtain D3-brane solutions which spread over EH-resolved conifold. The solutions are singular with the presence of 3-form flux background. Switching off the flux will make the solutions regular. However taking near horizon limit does not immediately give us $AdS_5 \times T^{1,1}$ space. The AdS geometry exists only for the branes located away from the deformed region of the cone.

While we were finishing this work two simultaneous works have appeared [16, 17] which also discuss AdS/CFT on the resolved S^2 and $S^2 \times S^2$ -conifolds. To add, the generalised Eguchi-Hanson solitons in odd dimensions have been constructed in [18].

Acknowledgements

I wish to thank S. Govindrajan for a fruitful discussion on conifold.

References

- [1] J.M. Maldacena, Adv. Theor. Math. Phys. 2 (1998) 231.

- [2] S.S. Gubser, I.R. Klebanov, A.M. Polyakov, Phys. Lett. 428B (1998) 105.
- [3] E. Witten, Adv. Theor. Math. Phys. 2 (1998) 253.
- [4] G. 't Hooft, gr-qc/9310006;
L. Susskind, J. Math. Phys. 36 (1995) 6377.
- [5] O. Aharony, S. Gubser, J. Maldacena, H. Ooguri, Y. Oz, Phys. Rept. 323 (2000) 183;
G.T. Horowitz and J. Polchinski, gr-qc/0602037.
- [6] P. Candelas and X. C. de la Ossa, *Comments on Conifolds*, Nucl. Phys. **B342** (1990) 246.
- [7] I.R. Klebanov and M.J. Strassler, *Supergravity and a Confining Gauge Theory: Duality Cascades and χ SB-Resolution of Naked Singularities*, hep-th/0007191.
- [8] L.A. Pando Zayas and A.A. Tseytlin, 3-branes on spaces with $R \times S^2 \times S^3$ topology, Phys. Rev. D **63** (2001) 086006, hep-th/0101043.
- [9] W. Chen, H. Lü, C.N. Pope, *Kerr-de Sitter black holes with NUT charges*, hep-th/0601002.
- [10] M. Bianchi, M.B. Green, S. Kovacs, G. Rossi, JHEP 9808 (1998) 013.
- [11] N. Dorey, V.V. Khoze, M.P. Mattis, S. Vandoren, Phys. Lett. B442 (1998) 145.
- [12] T. Eguchi and A.J. Hanson, Ann. Phys. 120 (1979) 82; T. Eguchi and A.J. Hanson, Phys. Lett. B74 (1978) 249.
- [13] R.R. Parwani and H. Singh, *Information geometry of $U(1)$ instantons on self-dual manifolds*, hep-th/0608144.
- [14] M. Blau, K.S. Narain, G. Thompson, hep-th/0108122.
- [15] I.R. Klebanov and E. Witten, *AdS/CFT Correspondence and Symmetry Breaking*, Nucl. Phys. **B556** (1999) 89, hep-th/9905104.
- [16] I.R. Klebanov and A. Murugan, *Gauge/Gravity Duality and Warped Resolved Conifold*, hep-th/0701064.
- [17] W. Chen, M. Cvetič, H. Lü, C.N. Pope and J.F. Vazquez-Poritz, *Resolved Calabi-Yau Cones and Flows from L^{abc} Superconformal Field Theories*, hep-th/0701082.
- [18] R. Clarkson and R. Mann, *Eguchi-Hanson Solitons in Odd Dimensions*, Class. Quant. Grav. 23:1507-1524 (2006); hep-th/0508200.